

SOLICARE ODT

5 mg and 10 mg orodispersible tablets

Contains the active ingredient solifenacin succinate

Consumer Medicine Information

What is in this leaflet

Read this leaflet carefully before you start taking this medicine.

This leaflet answers some common questions about SOLICARE ODT.

It does not contain all of the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have benefits and risks. Your doctor has weighed the risks of you taking SOLICARE ODT against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, talk to your doctor or pharmacist.

Keep this leaflet with your medicine. You may need to read it again.

What SOLICARE ODT is used for

SOLICARE ODT is used to treat the symptoms of overactive bladder. These symptoms include needing to rush to the toilet to urinate without warning, needing to urinate frequently, or wetting yourself because you did not get to the toilet in time.

SOLICARE ODT belongs to a group of medicines called anticholinergic medicines. It works by acting on specialised cells to reduce the activity of your bladder and help you control your bladder.

Your doctor may have prescribed SOLICARE ODT for another

reason. Ask your doctor if you have any questions about why SOLICARE ODT has been prescribed for you.

SOLICARE ODT is available only with a doctor's prescription.

There is no evidence that SOLICARE ODT is addictive.

Before you take SOLICARE ODT

When you must not take it

Do NOT take SOLICARE ODT if you are allergic to medicines containing solifenacin or any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include skin rash, itching or hives, swelling of the face, lips or tongue which may cause difficulty in swallowing or breathing, wheezing or shortness of breath.

Do not take SOLICARE ODT if you:

- suffer from urinary retention (inability to urinate or empty your bladder completely)
- are undergoing haemodialysis
- have severe kidney disease or moderate liver disease and are also taking medicines that can decrease the removal of SOLICARE ODT from the body (see Taking other medicines)
- have severe liver disease
- have uncontrolled glaucoma (high pressure in the eyes, with gradual loss of eye sight)

- have myasthenia gravis, a disease of the muscles causing drooping eyelids, double vision, difficulty in speaking or swallowing and sometimes muscle weakness in the arms or legs
- have a blocked gastro-intestinal tract or other gastro-intestinal disorders such as ulcerative colitis or toxic megacolon.

Do not take SOLICARE ODT if the expiry date (Exp.) printed on the pack has passed.

Do not take SOLICARE ODT if the packaging is torn or shows signs of tampering.

Before you start to take it

Tell your doctor if you are allergic to any other medicines, foods, dyes or preservatives.

Do not give SOLICARE ODT to children.

Tell your doctor if you are pregnant.

SOLICARE ODT may affect your developing baby if you take it during pregnancy. Your doctor will decide if you should take it.

Do not take SOLICARE ODT if you are breastfeeding.

Tell your doctor if you have, or have had, any medical conditions, especially the following:

- low potassium levels or a family history of heart rhythm problems. SOLICARE ODT may have an effect on the electro-cardiogram (ECG - a heart tracing) and may add to the effect of other medicines on the ECG. You should advise

your doctor of any other medicines you are currently taking, particularly those affecting the heart rhythm. Your doctor may wish to perform an ECG if you have any risks of having an abnormal heart rhythm.

- blockage to your bladder that causes difficulty urinating
- obstruction of the digestive system (constipation)
- glaucoma (high pressure in the eyes, with gradual loss of eye sight)
- hiatus hernia or gastric reflux or are currently being treated with some medicines (such as bisphosphonates) which may make your reflux worse
- autonomic neuropathy, a nerve problem which sometimes occurs with diabetes and can lead to diarrhoea, impotence or low blood pressure
- severe kidney disease
- moderate liver disease.

If you have not told your doctor about any of the above, tell him/her before you start taking SOLICARE ODT.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines may be affected by SOLICARE ODT, or may affect how well it works.

Some medicines can slow down the removal of SOLICARE ODT from the body. If you are taking any of the following, 5 mg a day is the maximum recommended dosage:

- some antifungal medicines such as ketoconazole, Itraconazole
- ritonavir and nelfinavir, medicines to treat HIV infection.

If you are taking any of the above medicines and you also have severe kidney disease or moderate

liver disease, you should not take SOLICARE ODT.

If you are not sure whether you are taking any of these medicines, check with your doctor or pharmacist.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking SOLICARE ODT.

How to take SOLICARE ODT

How much to take

The usual dose is 5 to 10 mg of SOLICARE ODT once a day. Your doctor will decide the most suitable dose for you.

Talk to your doctor or pharmacist if you feel that the effect of SOLICARE ODT is too strong or too weak.

Follow all directions given to you by your doctor and pharmacist carefully.

How to take SOLICARE ODT

Try and take SOLICARE ODT at the same time each day. SOLICARE ODT is taken by mouth.

SOLICARE ODT should be taken orally and should be sucked until completely disintegrated. It can be taken with or without food.

If you forget to take SOLICARE ODT

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take the missed dose as soon as you remember, and then go back to taking your tablets as you would normally.

Do not take a double dose to make up for the dose you missed.

If you are not sure what to do, ask your doctor or pharmacist.

How long to take SOLICARE ODT for

To properly control your condition, SOLICARE ODT must be taken every day.

Keep taking SOLICARE ODT for as long as your doctor recommends.

If you take too much SOLICARE ODT (overdose)

Immediately telephone your doctor, or the Poisons Information Centre (telephone 13 11 26) if you think you or anyone else may have taken too much SOLICARE ODT. Do this even if there are no signs of discomfort or poisoning.

While you are taking SOLICARE ODT

Things you must do

Before starting any new medicine, tell your doctor or pharmacist that you are taking SOLICARE ODT.

Tell all the doctors, dentists and pharmacists who are treating you that you are taking SOLICARE ODT.

If you become pregnant while taking SOLICARE ODT, tell your doctor.

Visit your doctor regularly so they can check on your progress.

Things you must not do

Do not stop taking SOLICARE ODT, or lower the dose, without checking with your doctor.

Do not use SOLICARE ODT to treat any other conditions unless your doctor tells you to.

Do not give SOLICARE ODT to anyone else, even if they have the same condition as you.

Things to be careful of

Be careful driving or operating machinery until you know how SOLICARE ODT affects you.

SOLICARE ODT can sometimes cause blurred vision and less frequently drowsiness and fatigue.

Do not drive or operate machinery if you suffer from any of these side effects.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking SOLICARE ODT.

Like all other medicines, SOLICARE ODT may have unwanted side effects in some people. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Stop taking SOLICARE ODT and tell your doctor as soon as possible if you experience:

- allergic reactions including swelling of the face, lips, throat or tongue, difficulty breathing

Tell your doctor as soon as possible if you notice any of the following:

- chest pain or tightness
- swelling of the feet or legs
- changes in electrical activity of the heart (ECG), irregular heartbeat (Torsade de Points)

Tell your doctor if you notice any of the following and they worry you:

- dry mouth
- nausea, vomiting
- constipation

- indigestion
- heartburn (dyspepsia)
- stomach discomfort
- urinary tract infection, bladder pain
- difficulty passing urine, inability to empty the bladder
- urgent need to urinate
- blurred vision
- inability to empty the bladder
- dry eyes
- tiredness
- muscle weakness
- accumulation of fluid (swelling) in the lower limbs (oedema)
- dizziness
- sore / dry throat
- cough
- headache
- sleepiness
- hallucination
- confusion
- depression
- fast heartbeat
- palpitations
- skin reactions such as rash, itching or hives.

Other side effects not listed above may also occur in some patients.

Tell your doctor if you notice anything that is making you feel unwell.

Some of these side effects (for example, changes in electrical activity of the heart or liver function) can only be found when your doctor does tests from time to time to check your progress.

After using SOLICARE ODT

Storage

Keep SOLICARE ODT where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Keep your tablets in a cool dry place where the temperature stays below 25°C.

Do not store SOLICARE ODT or any other medicine in the bathroom or near a sink.

Do not leave SOLICARE ODT in the car or on window sills.

Heat and dampness can destroy some medicines.

Disposal

If your doctor tells you to stop taking SOLICARE ODT, or your tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.

Product description

What it looks like

SOLICARE ODT comes in two strengths of tablets:

- SOLICARE ODT 5 mg orodispersible tablets are white, round, biconvex tablets, debossed with 5 on one side.
- SOLICARE ODT 10 mg orodispersible tablets are white, round, biconvex tablets.

Each blister pack contains 30 tablets.

Ingredients

The active ingredient in SOLICARE ODT is solifenacin succinate

- each SOLICARE ODT 5 tablet contains 5 mg solifenacin succinate
- each SOLICARE ODT 10 tablet contains 10 mg solifenacin succinate

The tablets also contain:

- polacrillin potassium
- hypromellose
- mannitol
- sucralose
- peppermint flavour
- lactose monohydrate

- croscarmellose sodium
- menthol aroma
- sodium stearyl fumarate

This medicine contains lactose and sucralose.

Sponsor:

Arrow Pharma Pty Ltd
15-17 Chapel Street
Cremorne VIC 3121
www.arrowpharma.com.au

This leaflet was prepared in
December 2021.

Australian registration numbers:

5 mg tablet: AUST R 316984

10 mg tablet: AUST R 316987