

CHEMISTS' OWN OSTEORELIEF PARACETAMOL

665 mg Paracetamol

Consumer Medicine Information

Please read this information before you start using this medicine.

What is in this leaflet

This leaflet answers some common questions about CHEMISTS' OWN OSTEORELIEF PARACETAMOL. It does not contain all the available information.

It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits.

If you have any concerns about using this medicine, ask your pharmacist or doctor.

Keep this leaflet with the medicine. You may need to read it again.

What CHEMISTS' OWN OSTEORELIEF PARACETAMOL is used for

CHEMISTS' OWN OSTEORELIEF PARACETAMOL is effective for the relief of persistent pain associated with:

- Osteoarthritis
- Muscle aches and pains such as backache

CHEMISTS' OWN OSTEORELIEF PARACETAMOL is also used for the fast effective temporary relief of pain and discomfort associated with:

- Headache
- Tension headache
- Period pain
- Toothache and pain after dental procedures

- Colds and flu

Paracetamol also reduces fever.

Paracetamol works to stop the pain messages from getting through to the brain. It also acts in the brain to reduce fever.

Ask your pharmacist or doctor if you have any questions about this medicine.

Your pharmacist or doctor may have recommended it for another reason.

Before you use CHEMISTS' OWN OSTEORELIEF PARACETAMOL

Do not take more than the recommended dose as it may cause serious harm to your liver.

When you must not use it

Do not use CHEMISTS' OWN OSTEORELIEF PARACETAMOL if you have an allergy to:

- Any medicine containing paracetamol
- Any of the ingredients listed at the end of this leaflet

Do not take this medicine if you are taking other prescription or non-prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu, or to aid sleep.

Always read and follow the label.

Do not use this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start using this medicine, talk to your pharmacist or doctor.

Before you start to use it

Tell your pharmacist or doctor if you have or have had any of the following medical conditions:

- Liver or kidney disease
- Are underweight or malnourished
- Regularly drink alcohol
You may need to avoid using this product altogether or limit the amount of paracetamol that you take.
- You have a severe infection, are severely malnourished or are a chronic heavy alcohol user as this may increase the risk of metabolic acidosis:

Signs of metabolic acidosis include:

- Deep, rapid, difficulty breathing
- Feeling sick (nausea), being sick (vomiting)
- Loss of appetite

Contact a doctor immediately if you get a combination of these symptoms.

Please see your doctor if symptoms do not improve.

Keep out of sight and reach of children.

Ask your pharmacist or doctor about using paracetamol if you are

pregnant or plan to become pregnant, or breastfeeding.

Paracetamol may be used during pregnancy and if you are breastfeeding, but you should always consult your doctor first.

Consider taking the lowest effective dose for the shortest period of time.

If you have not told your pharmacist or doctor about any of the above, tell them before you use CHEMISTS' OWN OSTEORELIEF PARACETAMOL.

Using other medicines

Tell your pharmacist or doctor if you are using any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and paracetamol may interfere with each other. These include:

- Warfarin, a medicine used to prevent blood clots
- Metoclopramide, a medicine used to control nausea and vomiting
- Medicines used to treat epilepsy or fits
- Chloramphenicol, an antibiotic used to treat ear and eye infections
- Alcohol
- Probenecid, a medicine used to treat gout or sometimes given with an antibiotic
- Cholestyramine, a medicine used to treat high cholesterol levels in the blood

Your pharmacist and doctor will have more information on these and other medicines to be careful with or avoid while using the medicine.

How to use CHEMISTS' OWN OSTEORELIEF PARACETAMOL

Follow all directions given to you by your pharmacist or doctor carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the box, ask your pharmacist or doctor for help.

Do not exceed the stated dose.

Use the smallest dose that you need to treat your symptoms and use the medicine for the shortest period of time necessary.

How much to use

Adults and children aged 12 years and over: Take 2 tablets three times a day, every six to eight hours as needed. Do not take more than 6 tablets in 24 hours.

Not recommended for children under 12 years.

How to use it

Swallow the tablets whole with water or other fluid.

Do not crush the tablets. They can be taken with or without food.

Try to space the doses at equal intervals throughout the day.

How long to use it

Adults and children aged 12 years and over: Only take paracetamol for a few days at a time unless your doctor tells you to take it for longer.

If you use too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much CHEMISTS' OWN OSTEORELIEF PARACETAMOL.

Do this even if there are no signs of discomfort or poisoning because of the risk of liver failure.

You may need urgent medical attention.

While you are using CHEMISTS' OWN OSTEORELIEF PARACETAMOL

Things you must do

Talk to your pharmacist or doctor if your symptoms do not improve. Your pharmacist or doctor will assess your condition and decide if you should continue to take the medicine.

Things you must not do

Do not take this medicine for longer than a few days at a time unless your doctor tells you to.

Do not take more than the recommended dose unless your doctor tells you to.

Do not use CHEMISTS' OWN OSTEORELIEF PARACETAMOL to treat any other complaints unless your pharmacist or doctor tells you to.

Only drink small quantities of alcohol (beer, wine or spirits) while using CHEMISTS' OWN OSTEORELIEF PARACETAMOL.

Drinking large quantities of alcohol while taking CHEMISTS' OWN OSTEORELIEF PARACETAMOL may increase the risk of liver side effects.

Side Effects

Tell your pharmacist or doctor as soon as possible if you do not feel well while using CHEMISTS' OWN OSTEORELIEF PARACETAMOL.

This medicine helps most people with various types of pain but it may have unwanted side effects. All medicines can have side effects. Sometimes they are serious, most of

the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following list of side effects. These side effects are rare and you may not experience any of them.

Ask your pharmacist or doctor to answer any questions you may have.

If any of the following happen, stop using the product and tell your pharmacist or doctor immediately or go to Accident and Emergency at your nearest hospital:

- Shortness of breath
- Wheezing or difficulty breathing
- Swelling of the face, lips, tongue or other parts of the body
- Rash, peeling, itching or hives on the skin or mouth ulcers
- Unexplained bruising or bleeding

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

These side effects are very rare for low doses of this medicine and when used for a short period of time.

Tell your pharmacist or doctor if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After using CHEMISTS' OWN OSTEO RELIEF PARACETAMOL

Storage

Keep your medicine in the original pack until it is time to take it.

Keep your medicine in a cool dry place where the temperature stays below 25°C.

Do not store CHEMISTS' OWN OSTEO RELIEF PARACETAMOL or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in a car.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

Ask your pharmacist what to do with any medicine that is left over, or if the expiry date has passed.

Product Description

What it looks like

CHEMISTS' OWN OSTEO RELIEF PARACETAMOL tablets are a white to off-white, film-coated capsule shaped tablet with flat edges. They are marked "8" on one side and are plain on the other side. They come in blister packs of 6 and 96 tablets.

Ingredients

CHEMISTS' OWN OSTEO RELIEF PARACETAMOL tablets contain 665 mg of paracetamol as the active ingredient.

They also contain:

- Colloidal anhydrous silica
- Hypromellose
- Macrogol 6000
- Magnesium stearate
- Maize starch
- Microcrystalline cellulose
- Povidone
- Pregelatinized maize starch
- Purified talc
- Sodium starch glycollate type A
- Titanium dioxide

Manufacturer/Supplier

CHEMISTS' OWN OSTEO RELIEF PARACETAMOL tablets are supplied in Australia by:

Arrotex Pharmaceuticals Pty Ltd
15 – 17 Chapel Street
Cremorne VIC 3121

AUST R 227111

Date of preparation: November 2022.