

Imatinib GH

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using Imatinib GH?

Imatinib GH contains the active ingredient imatinib mesilate. Imatinib GH is used to treat adults who have certain types of cancer and leukaemia.

For more information, see Section [1. Why am I using Imatinib GH ?](#) in the full CMI.

2. What should I know before I use IMATINIB GH?

Do not use if you have ever had an allergic reaction to Imatinib GH or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I use Imatinib GH?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with Imatinib GH and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use Imatinib GH?

- Your doctor will tell you exactly how many tablets of Imatinib GH to take
- Follow the instructions provided and use Imatinib GH until your doctor tells you to stop.

More instructions can be found in Section [4. How do I use Imatinib GH ?](#) in the full CMI.

5. What should I know while using Imatinib GH ?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using Imatinib GH.• Have regular check-ups to make sure the treatment is working
Things you should not do	<ul style="list-style-type: none">• Do not stop using this medicine suddenly (unless your doctor tells you to)
Driving or using machines	<ul style="list-style-type: none">• Be careful before you drive or use any machines or tools until you know how Imatinib GH affects you.
Drinking alcohol	<ul style="list-style-type: none">• There are no known interactions between Imatinib GH and alcohol
Looking after your medicine	<ul style="list-style-type: none">• Store it in a cool dry place away from moisture, heat or sunlight.• Keep out of reach of children.

For more information, see Section [5. What should I know while using Imatinib GH ?](#) in the full CMI.

6. Are there any side effects?

Common side effects include headache, dizziness, light-headedness or vertigo, upset stomach, nausea (feeling sick), vomiting, diarrhoea, constipation, rash, eczema, itchy or dry skin, skin darkening or lightening, symptoms of sunburn which happens more quickly than normal, noticeable swelling, stiffness in joints or muscles, pain in bones or along veins, weakness, feeling generally unwell, numbness, tingling in fingers and toes, change in sense of taste, irritated eyes, blurred vision, ringing in the ears, changes in appetite and weight, hair loss, throat pain, cough or cold symptoms, loss of interest in sex, problems with sexual function, breast enlargement, nipple pain, painful periods, slowing of growth in children and adolescents

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

Imatinib GH

Active ingredient(s): *Imatinib mesilate*

Consumer Medicine Information (CMI)

This leaflet provides important information about using Imatinib GH. You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Imatinib GH.

Where to find information in this leaflet:

- [1. Why am I using Imatinib GH ?](#)
- [2. What should I know before I use Imatinib GH ?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Imatinib GH ?](#)
- [5. What should I know while using Imatinib GH ?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using Imatinib GH ?

Imatinib GH contains the active ingredient Imatinib mesilate. Imatinib GH belongs to a group of anti-cancer medicines known as tyrosine kinase inhibitors.

Imatinib GH is used to treat adults who have certain types of cancer and leukemia, including chronic myeloid leukemia (CML) and acute lymphoblastic leukemia with Philadelphia chromosome positive (Ph-positive ALL).

Imatinib GH is also used to treat adults for:

- **Myelodysplastic / myeloproliferative diseases (MDS/MPD).** These are a group of blood diseases in which some blood cells start growing out of control.
- **Aggressive systemic mastocytosis (ASM).** It is a cancer in which certain blood cells, called "mast" cells, grow out of control.
- **Hypereosinophilic syndrome (HES) and or chronic eosinophilic leukaemia (CEL).** These are blood diseases in which some blood cells, named "eosinophils", start growing out of control.
- **Gastro-intestinal stromal tumours (GIST).** This is a type of cancer of the stomach and bowels. This cancer affects the tissue that surrounds the stomach and bowels and the cells grow uncontrollably.
- **Dermatofibrosarcoma protuberans (DFSP).** DFSP is a cancer of the tissue beneath the skin in which some cells start growing out of control.

Imatinib GH works by slowing the growth of and killing cancer cells while leaving normal cells alone.

2. What should I know before I use Imatinib GH ?

Warnings

Do not use Imatinib GH if:

- you are allergic to imatinib, or any of the ingredients listed at the end of this leaflet.

Tell your doctor if you are allergic to any other medicines, foods, dyes or preservatives.

If you experience an allergic reaction, stop using the medicine and inform your doctor or pharmacist immediately.

- Some of the symptoms of an allergic reaction may include
 - shortness of breath, wheezing or difficulty breathing;
 - swelling of the face, lips, tongue or other parts of the body;
 - rash, itching or hives on the skin.
- you have the rare hereditary problems of glucose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption as Cipla Imatinib contains lactose

Check with your doctor if you:

- have any other medical conditions
- take any medicines for any other condition

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

This medicine may be harmful to your unborn baby. If it is necessary for you to take it during pregnancy, your doctor will let you know.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

It is not known if the active ingredient, imatinib, passes into the breast milk. Breastfeeding is not recommended because this medicine could affect your baby.

Tell your doctor if you have ever had any of the following medical problems or procedures:

- kidney or liver problems
- problems with your heart
- you have had your thyroid gland removed
- hepatitis B infection. As during treatment with Imatinib GH, hepatitis B may become active again. Your doctor may want to take special precautions in this case.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interfere with Imatinib GH and affect how it works. These include many medicines that are eliminated from the body through the liver e.g.

- St. John's wort, a herbal medicine found in many products that you can buy without a prescription
- paracetamol, a medicine found in many common pain relievers and cold remedies (e.g. Panadol®, Panadeine®, Codral®, Tylenol®) which are known to be associated with liver toxicity.
- antibiotic medicines such as rifampicin, ketoconazole, erythromycin, clarithromycin, itraconazole
- antiviral medicines used to treat HIV/AIDS
- dexamethasone, a steroid medicine
- medicines for high cholesterol, such as simvastatin
- medicines used to treat epilepsy, such as phenytoin, carbamazepine, phenobarbitone
- warfarin, a medicine used to prevent blood clots
- some medicines used to treat mental disorders and depression
- some medicines used to treat high blood pressure and heart problems
- cyclosporine

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Imatinib GH .

4. How do I use Imatinib GH ?

How much to take

- Your doctor will tell you exactly how many tablets of Imatinib GH to take
- Do not exceed the recommended dose prescribed by your doctor
- Follow the instructions provided and use Imatinib GH until your doctor tells you to stop
- Your doctor may direct you to take a higher or lower dose or stop treatment depending on how your body responds to Imatinib GH.
- Imatinib GH is usually taken as a single dose each day. However, your doctor may want you to take them in two doses, one in the morning and one in the evening

When to take Imatinib GH

Take Imatinib GH every day; at the same time, with water and food for as long as your doctor recommends.

How to take Imatinib GH

- Take the medicine with a large glass of water and food. This will help avoid stomach irritation
- If you are unable to swallow the capsules
 - you can open the capsule and pour the powder

capsule contents into a glass of still water or apple juice (approximately 50 mL for 100 mg capsule or 200 mL for 400 mg capsule)

- Stir with a spoon to complete mixing of the capsule
- Drink immediately

If you forget to use Imatinib GH

Imatinib GH should be taken regularly at the same time each day. If you miss your dose at the usual time, take the missed dose as soon as you remember, then continue with your normal schedule

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose you missed.

If you take too much Imatinib GH

If you think that you have taken too much Imatinib GH , you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while using Imatinib GH ?

Things you should do

Make sure you follow your doctor's instructions carefully and keep all appointments.

You will need regular check-ups to make sure the treatment is working. Regular blood tests, weight checks and urine tests can also find side effects before they become serious.

Use a method of contraception to prevent pregnancy during treatment and for 15 days after ending treatment.

Some children taking Imatinib GH may have slower than normal growth. Growth will be monitored at regular visits by your doctor

Call your doctor straight away if you:

- become pregnant or think you might be pregnant while you are taking this medicine.
- Think you might be having an allergic reaction to Imatinib GH treatment

Remind any doctor, dentist or pharmacist you visit that you are using Imatinib GH .

Things you should not do

- Do not stop using this medicine suddenly unless your doctor tells you to.

- Do not give this medicine to anyone else even if they have the same condition as you.

Drinking grapefruit juice

Avoid drinking grapefruit juice while you are being treated with Imatinib GH. Grapefruit juice may interact with Imatinib GH and affect how your body uses this medicine.

Treating a headache, cold or other minor aches and Pains

Try to avoid taking medicines containing paracetamol (e.g. Panadol®, Panadeine®, Codral®, Tylenol®) to treat minor aches and pains. Ask your pharmacist to suggest an alternative medicine.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how Imatinib GH affects you.

This medicine may cause dizziness, light-headedness or drowsiness in some people.

Wear sun protection when outdoors

When you are outdoors, wear protective clothing and use at least a 15+ sunscreen. Do not use a sunlamp or tanning beds.

This medicine may cause your skin to be much more sensitive to sunlight than it normally is.

Exposure to sunlight may cause a skin rash, itching, redness or severe sunburn. If your skin does appear to be burning, tell your doctor.

Drinking alcohol

Tell your doctor if you drink alcohol.

There are no known interactions of Imatinib GH with alcohol.

Looking after your medicine

- Keep your medicine in the original container until it is time to take it.

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary.

However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<p>Stomach and intestine problems:</p> <ul style="list-style-type: none"> • pain or burning sensation in upper tummy area and/or chest (heartburn), feeling sick, vomiting, feeling of fullness and bloating, black-coloured stool (signs of stomach ulcer). <p>Skin problems:</p> <ul style="list-style-type: none"> • skin rash with flaking or peeling (exfoliative dermatitis). • blisters on skin (signs of pemphigus). • painful and/or blistering skin areas • decreased or increased skin sensitivity. • changes in skin colour. • red bumps or white-headed pimples around the roots of the hair, possibly with pain, itching or burning feeling (signs of swelling of the hair follicles, also called folliculitis). • painful red lumps on the skin, skin pain, skin reddening (inflammation of fatty tissue under the skin, also called panniculitis). <p>General problems:</p> <ul style="list-style-type: none"> • feeling dizzy or weak • itching. • unusual hair loss or thinning • dry mouth, dry skin or dry eye • hot flushes, chills or night sweats • severe headache felt as a throbbing pain or pulsing sensation, usually on one side of the head and often with sick feeling, vomiting and sensitivity to light or sound (signs of migraine). • overwhelming urge to move the legs (restless leg syndrome). • burping/belching. • difficulty swallowing • increased sweating • brittle nails • generally feeling unwell • increase in appetite. • nail discolouration. • slowing of growth in children and adolescents. 	Speak to your doctor if you have any of these less serious side effects and they worry you.

<p>Ear problems:</p> <ul style="list-style-type: none"> hearing noises (e.g. ringing, humming) in the ears that have no external source (tinnitus). <p>Eye problems:</p> <ul style="list-style-type: none"> discharge from the eye with itching, redness and swelling (conjunctivitis), watery eyes or having blurred vision. <p>Blood problems:</p> <ul style="list-style-type: none"> nose bleeds. <p>Infection problems:</p> <ul style="list-style-type: none"> viral infection such as cold sore. swelling of the lips. mouth ulcers. cough, runny or stuffy nose, feeling of heaviness or pain on pressing the area above the eyes or on the sides of the nose, nasal stuffiness, sneezing, sore throat, with or without headache (signs of upper respiratory tract infection). flu-like symptoms (influenza). pain or burning feeling while passing urine, increased body temperature, pain in groin or pelvic area, red-or brown-coloured or cloudy urine (signs of urinary tract infection). <p>Nutrition problems:</p> <ul style="list-style-type: none"> anorexia, weight loss or a change in sense of taste. <p>Bone and connective tissue problems:</p> <ul style="list-style-type: none"> joint pain with swelling. pain and swelling of your joints (signs of arthralgia). <p>Kidney problems:</p> <ul style="list-style-type: none"> Increased frequency of passing urine lower back pain resulting from kidney problems <p>Muscle problems:</p> <ul style="list-style-type: none"> joint and muscle stiffness. <p>Nervous system problems:</p> <ul style="list-style-type: none"> difficulty in sleeping (insomnia). a constant feeling of sadness and loss of interest, which stops you carrying out your normal activities (signs of depression). a feeling of nervousness and worry along with physical changes such as pounding heart, sweating, 	<p>trembling, dry mouth (signs of anxiety).</p> <ul style="list-style-type: none"> sleepiness/drowsiness/excessive sleep. confusion. memory loss. trembling or shaky movements (tremor). <p>Heart and circulation problems:</p> <ul style="list-style-type: none"> numbness of the hands or feet. reddening and/or swelling on the palms of the hands and soles of the feet with tingling feeling and burning pain. <p>Reproduction problems:</p> <ul style="list-style-type: none"> breast enlargement (may occur in men or women). dull pain and/or feeling of heaviness in the testicles or lower tummy area, pain during pee, sexual intercourse or ejaculation, blood in urine (signs of oedema of the testicles). unable to get or keep an erection (erectile dysfunction). heavy or irregular menstrual periods. difficulty achieving/maintaining sexual arousal. decreased sexual desire. nipple pain. slowing of growth in children and adolescents.
---	---

Serious side effects

Serious side effects	What to do
<p>Blood problems:</p> <ul style="list-style-type: none"> unexpected bleeding or bruising (when you have not hurt yourself). feeling light-headed, dizzy or fainting (signs of low blood pressure). pale skin, feeling tired, difficulty breathing or breathless and having dark urine (signs of low levels of red blood cells). numb or cold toes and fingers (signs of Raynaud's syndrome). muscle weakness and spasms with an abnormal heart rhythm (signs of changes in the amount of potassium in your blood). bruising. feeling sick, shortness of breath, irregular heartbeat, cloudy urine, tiredness and/or joint pain with 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

<p>abnormal blood test results (eg. high potassium, uric acid and calcium levels and low phosphorous levels in the blood).</p> <ul style="list-style-type: none"> tired, dizziness, shortness of breath, bruises, gum/nose bleeds, minor cuts bleed a lot, feeling confused, sleepiness, seizures, decreased urine, swollen legs, fever (problems with blood clotting). <p>Stomach and intestine problems:</p> <ul style="list-style-type: none"> severe stomach pain, blood in vomit, stool or urine, black stools (signs of gastrointestinal problems) stomach pain with feeling sick (nausea) feeling sick (nausea), being sick (vomiting), diarrhoea, or upset stomach or heartburn <p>General problems:</p> <ul style="list-style-type: none"> rapid weight gain, Imatinib GH may cause your body to keep water (severe fluid retention). combined widespread severe rash, feeling sick, fever, high level of certain white blood cells or yellow skin or eyes (signs of jaundice) with difficulty breathing, chest pain, severely decreased urine output and feeling thirsty etc. (signs of a treatment-related allergic reaction). headache or feeling tired. swelling such as around your ankles or puffy eyes. weight gain. <p>Skin problems:</p> <ul style="list-style-type: none"> rash rash, red skin with blisters on the lips, eyes, skin or mouth, peeling skin, fever, raised red or purple skin patches, itching, burning feeling (signs of skin problems) sudden swelling and redness of the skin (signs of a skin infection called cellulitis). <p>Ear problems:</p> <ul style="list-style-type: none"> difficulty hearing <p>Eye problems:</p> <ul style="list-style-type: none"> eye pain or worsening eyesight, bleeding in the eyes <p>Liver problems:</p> <ul style="list-style-type: none"> Feeling sick (nausea), with loss of appetite, dark-coloured urine, 	<p>yellow skin or eyes (signs of liver problems).</p> <p>Infection problems:</p> <ul style="list-style-type: none"> Signs of infection such as fever, severe chills, sore throat or mouth ulcers. Imatinib GH can reduce the number of white blood cells, so you might get infections more easily. <p>Bone and connective tissue problems:</p> <ul style="list-style-type: none"> muscle cramps or joint, muscle or bone pain, during Imatinib GH treatment or after you have stopped taking Imatinib GH. pain in bones or joints (signs of osteonecrosis). <p>Nervous system problems:</p> <ul style="list-style-type: none"> severe headache, weakness or paralysis of arms/legs or face, difficulty speaking, sudden loss of consciousness (signs of nervous system problems such as bleeding or swelling in skull/brain). <p>Heart problems:</p> <ul style="list-style-type: none"> chest pain, irregular heart rhythm (signs of heart problems). <p>Excretory system problems:</p> <ul style="list-style-type: none"> feeling sick (nausea) with diarrhoea and vomiting, stomach pain or fever (signs of bowel problems). <p>Muscle problems:</p> <ul style="list-style-type: none"> muscle spasms with a fever, red-brown urine, pain or weakness in your muscles.
--	---

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Some of these side effects can only be found by laboratory testing.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What Imatinib GH contains

Active ingredient (main ingredient)	imatinib mesilate
Other ingredients (inactive ingredients)	<p><u>The other ingredients are:</u></p> <ul style="list-style-type: none">• lactose,• crospovidone,• colloidal anhydrous silica,• magnesium stearate. <p><u>The capsule shells (cap and body) contain:</u></p> <ul style="list-style-type: none">• gelatin,• purified water,• iron oxide red• iron oxide yellow• titanium dioxide.• sodium lauryl sulfate

Do not take this medicine if you are allergic to any of these ingredients.

What Imatinib GH looks like

Imatinib GH 100 mg capsules are light yellow granules filled in size 3 hard gelatin capsule with brown cap and white body (AUST R 281535).

Imatinib GH 400 mg capsules are light yellow granules filled in size 00EL hard gelatin capsules with brown cap and brown body (AUST R 281536).

Who distributes Imatinib GH

Generic Health Pty Ltd
Level 1, 1102 Toorak Road
CAMBERWELL VIC 3124
drugsafety@cipla.com
Phone no: 1 800 569 074

This leaflet was prepared in August 2022.