

Panadol® SINUS RELIEF

ORIGINAL FORMULA

Paracetamol 500 mg and Pseudoephedrine Hydrochloride 30 mg

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about PANADOL SINUS RELIEF ORIGINAL FORMULA.

It does not contain all the available information.

It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits. Your pharmacist or doctor has weighed the risks of you taking PANADOL SINUS RELIEF ORIGINAL FORMULA against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your pharmacist or doctor.

Keep this leaflet with the medicine.

You may need to read it again.

What PANADOL SINUS RELIEF ORIGINAL FORMULA is used for

PANADOL SINUS RELIEF ORIGINAL FORMULA is used for the temporary relief of sinus congestion and pain, nasal congestion and runny nose.

PANADOL SINUS RELIEF ORIGINAL FORMULA caplets® contain the active ingredients paracetamol and pseudoephedrine hydrochloride.

Paracetamol works to stop the pain messages from getting through to

the brain. It also acts in the brain to reduce fever.

Pseudoephedrine hydrochloride belongs to a group of medicines called sympathomimetic decongestants.

It works by reducing congestion in the upper respiratory tract, including the nose, nasal passages and sinuses, and making it easier to breathe.

Ask your pharmacist or doctor if you have any questions about this medicine.

Your pharmacist or doctor may have given it for another reason.

This medicine is not addictive.

It is only available from your pharmacist.

Before you take/give PANADOL SINUS RELIEF ORIGINAL FORMULA

Taking too much paracetamol can cause serious harm to your liver.

When you must not take it

Do not take PANADOL SINUS RELIEF ORIGINAL FORMULA if you have an allergy to:

- any medicine containing paracetamol or pseudoephedrine hydrochloride
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- shortness of breath

- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

Do not take this medicine if you:

- have high blood pressure
- have heart disease or problems with your blood vessels
- have kidney disease unless your doctor says you can
- have taken monoamine oxidase inhibitors, medicines used to treat depression, in the last 14 days
- are taking other medicines such as decongestants, appetite suppressants and amphetamine-like psychostimulants.

Do not give to children under 12 years of age.

Do not take this medicine if you are pregnant or plan to become pregnant.

It may affect your developing baby if you take it during pregnancy. Your pharmacist or doctor will discuss the benefits and possible risks of taking the medicine during pregnancy.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your pharmacist or doctor.

Before you start to take it

Tell your pharmacist or doctor if you have allergies to any other

medicines, foods, preservatives or dyes.

Tell your pharmacist or doctor if you have or have had any of the following medical conditions:

- liver disease
- kidney disease
- heart problems
- high blood pressure
- overactive thyroid gland
- diabetes
- heart disease and poor blood flow in the blood vessels of the heart
- glaucoma (high pressure in the eyes)
- prostate problems
- irregular heart beat
- a rare tumour of the adrenal gland (phaeochromocytoma)
- occlusive vascular disease (poor blood flow or narrowing of the blood vessels e.g. Raynaud's Phenomenon)
- Epilepsy
- asthma, bronchitis, chronic lung disorder, or other respiratory problems
- are underweight or malnourished
- regularly drink alcohol
You may need to avoid using this product altogether or limit the amount of paracetamol that you take
- a severe infection as this may increase the risk of metabolic acidosis
Signs of metabolic acidosis include:
 - deep, rapid, difficult breathing
 - feeling sick (nausea), being sick (vomiting)
 - loss of appetite

Contact a doctor immediately if you get a combination of these symptoms

Ask your pharmacist or doctor about taking PANADOL SINUS RELIEF ORIGINAL FORMULA if you are pregnant or plan to become pregnant, or breastfeeding.

Your pharmacist or doctor will discuss the potential benefits and risks of taking the medicine if you are pregnant or breastfeeding.

If you have not told your pharmacist or doctor about any of the above, tell him/her before you start taking PANADOL SINUS RELIEF ORIGINAL FORMULA.

Taking other medicines

Tell your pharmacist or doctor if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and PANADOL SINUS RELIEF ORIGINAL FORMULA may interfere with each other. These include:

- warfarin, a medicine used to prevent blood clots or similar medicines used to thin the blood
- pain relievers or medicines to treat anxiety, or help you sleep or relax (sedatives and hypnotics)
- metoclopramide, a medicine used to control nausea and vomiting
- anti-convulsants
- medicines used to treat epilepsy or fits such as phenytoin
- chloramphenicol, an antibiotic used to treat ear and eye infections
- alcohol
- medicines used to treat depression such as monoamine oxidase inhibitors (MAOIs)
- medicines used to treat heart conditions such as digoxin or similar medicines
- medicines used to treat high blood pressure such as beta-blockers

- medicines used to treat urinary tract infections and bladder problems
- medicines used to treat behavioural disorders
- phenylephrine, a medicine used to treat congestion
- appetite suppressants
- other decongestants or cough and cold medicines

These medicines may be affected by PANADOL SINUS RELIEF ORIGINAL FORMULA or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist will have more information on these and other medicines to be careful with or avoid while taking this medicine.

How to take PANADOL SINUS RELIEF ORIGINAL FORMULA

Follow all directions given to you by your pharmacist or doctor carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the box, ask your pharmacist or doctor for help.

Do not exceed the stated dose or frequency of dosing.

Always use the lowest effective dose to relieve your symptoms

How much to take

Adults and children 12 years and over: Take 2 caplets. Do not take more than 8 caplets in 24 hours.

Do not take more than the recommended dose.

How to take it

Take 2 caplets of PANADOL SINUS RELIEF ORIGINAL FORMULA by mouth with water.

When to take it

Every 6 hours as necessary

Maximum 8 caplets in 24 hours.

How long to take it

Adults:

Only take PANADOL SINUS RELIEF ORIGINAL FORMULA for a few days at a time except on medical advice.

Children aged 12 to 17 years:

Do not take for more than 48 hours except on medical advice.

Do not use in children below 12 years of age.

If you forget to take it

If it is less than 1 hour before your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember, and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your pharmacist or doctor.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much PANADOL SINUS RELIEF ORIGINAL FORMULA.

Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

While you are using PANADOL SINUS RELIEF ORIGINAL FORMULA

Things you must do

Talk to your pharmacist or doctor if your symptoms do not improve.

Your pharmacist or doctor will assess your condition and decide if you should continue to take the medicine.

Things you must not do

Children 12 to 17 years

Do not give PANADOL SINUS RELIEF ORIGINAL FORMULA for more than 48 hours unless a doctor has told you to.

Adults:

Do not take for more than a few days at a time unless your doctor tells you to.

Do not give to children under 12 years of age.

Do not take more than the recommended dose unless your doctor tells you to.

Do not take PANADOL SINUS RELIEF ORIGINAL FORMULA to treat any other complaints unless your pharmacist or doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Things to be careful of

Avoid drinking alcohol (beer, wine or spirits) while taking PANADOL SINUS RELIEF ORIGINAL FORMULA.

Drinking alcohol while taking PANADOL SINUS RELIEF ORIGINAL FORMULA may increase the risk of liver side effects.

Be careful driving or operating machinery until you know how PANADOL SINUS RELIEF ORIGINAL FORMULA affects you.

This medicine may cause dizziness in some people. If this happens, do not drive, use or operate machinery.

Side effects

Tell your pharmacist or doctor as soon as possible if you do not feel well while you are taking PANADOL SINUS RELIEF ORIGINAL FORMULA.

This medicine helps most people with the relief of sinus congestion and pain, nasal congestion and runny nose but there may be unwanted side effects in a few people.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Ask your pharmacist or doctor to answer any questions you may have.

Tell your pharmacist or doctor if you notice any of the following and they worry you:

- nausea or dyspepsia
- drowsiness
- difficulty sleeping
- nervousness
- agitation
- restlessness
- dizziness
- fear or anxiety
- rapid heart beat
- tremor
- increased blood pressure
- hallucinations
- trouble urinating (more likely to occur if you have enlarged prostate)

If any of the following happen, stop using the product, tell your pharmacist or doctor immediately or go to Accident and Emergency at your nearest hospital:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin
- unexplained bruising or bleeding
- rapid or irregular heartbeat
- urinary retention
- severe headache
- sudden abdominal pain or blood in your stools

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are very rare for low doses of this medicine and when used for a short period of time.

Tell your pharmacist or doctor if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After using PANADOL SINUS RELIEF ORIGINAL FORMULA

Storage

Keep your medicine in the original pack until it is time to take it.

Keep your medicine in a cool dry place where the temperature stays below 30°C.

Do not store PANADOL SINUS RELIEF ORIGINAL FORMULA or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

Ask your pharmacist what to do with any medicine that is left over, or if the expiry date has passed.

Product description

What it looks like

PANADOL SINUS RELIEF ORIGINAL FORMULA are white, capsule shaped tablets with flat edges debossed with I-I on one side.

A pack contains 24 tablets.

Ingredients

PANADOL SINUS RELIEF ORIGINAL FORMULA contains 500 mg of Paracetamol and 30 mg of Pseudoephedrine Hydrochloride as the active ingredients.

It also contains:

- Talc - purified
- Starch - Maize
- Starch - pregelatinised maize
- Stearic acid
- Povidone
- Sodium benzoate

Manufacturer/Supplier

PANADOL SINUS RELIEF ORIGINAL FORMULA is manufactured and supplied in Australia by:

Haleon Australia Pty Ltd
ACN 008 399 415

The Australian Registration number is:

AUST R 35521

Date of preparation: March 2023

Trademarks are owned by or licensed to the Haleon Group of Companies.